

# BIRTH AND BEYOND PEDIATRICS

## Advil or Motrin (Ibuprofen) Dosing Chart

Advil or Motrin		Infant Drops (50mg/1.25ml)	Children's Suspension (100mg/5ml)	Junior Strength Chewable tablets (100mg)	Junior Strength Caplets (100mg)
Weight (lbs)	Age				
13-17	6-11 mo.	1.25 mL		NA	NA
18-23	12-23 mo.	1.875 mL		NA	NA
24-35	2-3 yr	NA	1 tsp	1 tablet	NA
36-47	4-5 yr.	NA	1 ½ tsp	1 ½ tablets	NA
48-59	6-8 yr.	NA	2 tsp	2 tablets	2 tablets
60-71	9-10 yr.	NA	2 ½ tsp	2 ½ tablets	2 ½ tablets
72-95	11 yr.	NA	3 tsp	3 tablets	3 tablets

**For Advil or Motrin, give one dose every 6 to 8 hours.**

**Do not give to patients < 6mos. of age.**

**Do not give to patients with allergies to aspirin.**

## Tylenol (Acetaminophen) Dosing Chart

Tylenol		Infants' Drops (80mg/0.8ml)	Children's Suspension (160mg/5ml)	Children's Meltaway Tablets (80mg each)	Junior Tylenol Meltaway Tablets (160mg each)
		<b>OLD CONCENTRATION BEING DISCONTINUED</b>	<b>NEW FORMULA USED FOR INFANTS NOW</b>		
Weight (lbs)	Age				
6-11	0-3 mo.	0.4 mL	1.25 mL	NA	NA
12-17	4-11 mo.	0.8 mL	2.5 mL	NA	NA
18-23	12-23 mo.	1.2 mL	3.75 mL	NA	NA
24-35	2-3 yr.	1.6 mL	5 mL or 1 tsp.	2 tablets	NA
36-47	4-5 yr.	NA	7.5 mL or 1 ½ tsp.	3 tablets	NA
48-59	6-8 yr.	NA	10 mL or 2 tsp.	4 tablets	2 tablets
60-71	9-10 yr.	NA	12.5 mL or 2 ½ tsp.	5 tablets	2 ½ tablets
72-95	11 yr.	NA	15 mL or 3 tsp.	6 tablets	3 tablets
96 and up	12 yr.	NA	NA	NA	4 tablets

**For Tylenol, give one dose every 4 hours**

Infants' Acetaminophen Concentrated Dropper  
in 0.4 mL (40 mg) increments.

Infants' Acetaminophen Oral Suspension  
Syringe in 1.25 mL (40 mg) increments.